

**Table 1. Health Effect Levels of Temephos in Humans and Laboratory Animals**

Route	Duration	Species	NOAEL	LOAEL	Organ/Effect	Comments	Reference
<b>Acute Duration Toxicity</b>							
dermal	once	human	1.1 g/person		Signs of toxicity	Used from a shaker or duster in topical treatment for lice. Concluded to be safe and effective.	Steinberg et al. 1972
dermal	once	rat (m)		>4,000 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	INCHEM 2002
dermal	once	rat (f)		>4,000 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	INCHEM 2002
dermal	once	dog		>5,000 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	EXTOXNET 2002
dermal	once	cat		>5,000 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	EXTOXNET 2002
dermal	once	rabbit		1,300 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	INCHEM 2002
dermal	5 days	rabbit		0.4 ml/kg/day (178 mg a.i./kg/day)	Cholinesterase inhibition; diarrhea	Both cholinesterase inhibition and diarrhea were noted.	INCHEM 2002
oral	5 days	human	256 mg/day (~3.7 mg/kg/day)		Cholinesterase inhibition; clinical symptoms	No inhibition or symptoms observed.	Laws et al. 1967
oral	once	rat (m)		8,600 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	INCHEM 2002
oral	once	rat (f)		1,300 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	INCHEM 2002
oral	once	mouse		4,700 mg/kg	LD <sub>50</sub>	LD <sub>50</sub>	EXTOXNET 2002
oral	5 days	rabbit		100 mg/kg/day	Liver	Focal and diffuse hepatic necrosis noted.	HSDB 2003
oral	5 days	guinea pig	100 mg/kg/day		Organophosphate poisoning	No poisoning noted.	INCHEM 2002

**Table 1. Health Effect Levels of Temephos in Humans and Laboratory Animals (continued)**

Route	Duration	Species	NOAEL	LOAEL	Organ/Effect	Comments	Reference
<b>Intermediate Duration Toxicity</b>							
dermal	3 weeks, 5 days/wk	rat	12 mg/kg/day	60 mg/kg/day	Lesions; tissue changes; body weight	When applied dermally as aqueous emulsion, half of rats had abraded skin. Decreased weight gain noted in 60 mg/kg/day group (intact and abraded skin), but no other effects seen.	INCHEM 2002
oral	28 days	human	64 mg/day (~0.9 mg/kg/day)		Cholinesterase inhibition; clinical symptoms	No inhibition or symptoms observed.	Laws et al. 1967
oral	30 days	rabbit		10 mg/kg/day	Liver	Mild hepatic pathologic changes.	HSDB 2003
oral	44 day	rat	1 mg/kg/day	10 mg/kg/day	Erythrocyte cholinesterase inhibition	10 mg/kg/day resulted in 31% inhibition at 14 days and 47% at 44 days. No signs of organophosphate poisoning seen. Rats receiving 100 mg/kg/day showed signs of poisoning after 3 days (at 64% inhibition); gradual recovery from symptoms ensued, although inhibition progressed to 87% after 11 days.	INCHEM 2002
oral	35 days	rabbit	1 mg/kg/day	10 mg/kg/day	Cholinesterase inhibition; liver effects	No effects or significant inhibition were noted at 0.1 mg/kg/day or 1 mg/kg/day. The 10 mg/kg/day group developed 26% inhibition by day 7 and 47% inhibition by day 35. No animals showed signs of poisoning; no higher doses were used.	INCHEM 2002; HSDB 2003
oral	90 days	rat	6 ppm (0.3 mg/kg/day)	350 ppm (17.5 mg/kg/day)	Cholinesterase inhibition; clinical signs	Cholinesterase inhibition was the only effect noted.	INCHEM 2002; HSDB 2003
oral	35 days	rat	1 mg/kg/day		Cholinesterase inhibition	No cholinesterase inhibition was seen.	HSDB 2003
oral	186 days	sheep	5 mg/kg/day			No effects were noted.	HSDB 2003

**Table 1. Health Effect Levels of Temephos in Humans and Laboratory Animals (continued)**

Route	Duration	Species	NOAEL	LOAEL	Organ/Effect	Comments	Reference
oral	99 days	rat		2,000 ppm (~100 mg/kg/day)	Death, clinical signs, and erythrocyte cholinesterase inhibition	8 of 10 died within 10 days; 100% erythrocyte cholinesterase inhibition; signs of poisoning.	HSDB 2003
<b>Chronic Duration Toxicity</b>							
oral (in drinking water)	19 months	human	0.5 ppm		Cholinesterase inhibition; clinical symptoms	No inhibition or symptoms observed in village of 2,000 when water containers were treated once a month with temephos; 0.5 ppm estimated maximum dose.	Laws et al. 1968
oral (diet)	2 years	rat	300 ppm		Not specified	No effects were noted.	HSDB 2003
oral	422 days	sheep	2.5 mg/kg/day		Not specified	No effects were noted.	HSDB 2003
oral	1 year	cow		1 mg/kg/day	Not specified	“Signs of poisoning” were noted.	HSDB 2003
oral	2 years	rat	1 ppm	10 ppm	Liver effects	Minor pathologic changes noted in liver.	EXTOXNET 2002
<b>Developmental/Reproductive Toxicity</b>							
oral (in drinking water)	19 months	human	0.5 ppm		Cholinesterase inhibition; clinical symptoms; reproduction	No symptoms observed in village of 2,000 when water containers were treated once a month with temephos; 0.5 ppm estimated max. dose. Eight normal births were observed.	Laws et al. 1968
oral (diet)	3 generations	rat	125 ppm		Fertility, gestation, reproduction, lactation, congenital defects	No effects were noted. Dietary exposure continued from weaning through reproductive age.	HSDB 2003

**Table 1. Health Effect Levels of Temephos in Humans and Laboratory Animals (continued)**

<b>Route</b>	<b>Duration</b>	<b>Species</b>	<b>NOAEL</b>	<b>LOAEL</b>	<b>Organ/Effect</b>	<b>Comments</b>	<b>Reference</b>
oral	unspecified	rat		500 ppm (25 mg/kg/day)	Number of litters, litter size, viability, congenital defects, cholinesterase inhibition; signs of toxicity	No developmental or reproductive effects were noted, but some cholinesterase inhibition and toxicity were seen.	EXTOXNET 2002; HSDB 2003
oral	1 year	cow		1 mg/kg/day	Fertility	Evidence indicated that it may affect the fertility in heifers.	HSDB 2003